Large-eddy simulations of the response of tropopause moisture to deep convection

A. Ackerman, A. Fridlind, E. Jensen, A. Strawa NASA Ames Research Center

Dave Stevens LLNL

Larry Miloshevich NCAR

Paul Lawson SPEC, Inc.


Motivation


- Stratospheric moisture has increased over past decades (Oltmans and Hofmann, 1995)
- Stratospheric moisture plays important role in ozone depletion
- How does moisture enter the stratosphere?
 - 1. Gradual ascent through TTL
 - Convective overshooting (Sherwood and Dessler, 2000, 2001)


Approach: LES of deep convection over Florida


- LES (Stevens and Bretherton, 1997): $\Delta z=375$ m, $\Delta x=\Delta y=500$ m
- Explicit, bin microphysics model of aerosols, water, and ice (Toon, Turco, Jensen, Ackerman, Fridlind)
- Simulate three "golden" anvil days: July 11, 16, and 29
- Model domain initialized to observed, pre-convective sounding (massaged by L. Miloshevich)
- Open lateral boundaries, nudge to initial sounding with zeroed horizontal winds
- Initiate convection with warm bubble that reaches tropopause
- Aerosol concentrations: 500 cm⁻³ to 4 km, 150 cm⁻³ above


Case study: July 11
































































































!!! NOTE !!!

In the conversion from water vapor mass mixing ratio (as used by the model) to volume mixing ratio (as shown in the following plots) a mistake was made that resulted in the humidities being a factor of 4 smaller (as shown in the Utah presentation) than was actually measured.


This was only a plotting error -- the correct values were used in the model simulations.

The error has been corrected in this version of the plots.


July_11_Miami




July_11_PARSL




July_16_PARSL




July_29_Miami





July_29_PARSL





Isosurface of Condensed Water











Summary

- Simulations provide no evidence for dehydration of tropopause layer by deep convection
 - → Ice is lofted and evaporates faster than sedimentation can remove it in convective overshoots
- Significant disagreement in upper–level moisture between Miami (RS80–H) and PARSL (RS90) sondes
 - → Instrument artifact or real spatial variation?

Next steps

- Resolve radiosonde puzzle with in situ measurements of water vapor
- Explore sensitivities by varying aerosol concentrations, bubble strength, and initial soundings
- Analyze competing time scales (transport versus evaporation)
- Run cases from deep tropics mission